



IOWA DEPARTMENT OF NATURAL RESOURCES

---

# Iowa's 2022

# Forest Health Highlights

---



**December 2022**

**Jeff Goerndt, State Forester**

**Tivon Feeley, Forest Health Program Leader**

**Mark Runkel, Forest Health Program Technician**

# Contents

Introduction .....	1
Weather Review.....	1
Land Characteristics .....	4
United States Forest Service Major Pests List.....	5
United States Forest Service Major Pests List: Armillaria Root Disease .....	6
United States Forest Service Major Pests List: Asian long-horned beetle.....	7
United States Forest Service Major Pests List: Bur Oak Blight.....	8
United States Forest Service Major Pests List: Butternut Canker.....	10
United States Forest Service Major Pests List: Emerald Ash Borer .....	11
United States Forest Service Major Pests List: Forest Tent Caterpillar .....	13
United States Forest Service Major Pests List: <i>Lymantria dispar</i> .....	14
United States Forest Service Major Pests List: Heterobasidion Root Disease.....	17
United States Forest Service Major Pests List: Oak Wilt.....	18
United States Forest Service Major Pests List: Sudden Oak Death.....	19
United States Forest Service Major Pests List: Thousand Cankers Disease.....	20
United States Forest Service Major Pests List: Blister Rust .....	24
Additional Pest Surveyed: Spotted Lanternfly .....	25
Additional Pest Surveyed: Cankerworm .....	26
Additional Pest Surveyed: Pine Shoot Beetle .....	27
Additional Pest Surveyed: Dutch Elm Disease .....	29
Additional Pest Surveyed: Hickory Dieback .....	30
Additional Pest Surveyed: Invasive Plants .....	31
Additional Pest Surveyed: White Oak Mortality .....	34
Additional Pest Surveyed: Oak Tatters .....	35
Conclusion.....	36
Useful Phone Numbers and Websites .....	37

This project was funded in part through a grant awarded by the USDA, Forest Service, Northeastern Area State and Private Forestry.

Federal and state regulations prohibit discrimination on the basis of race, color, national origin, sex or disability. State law also prohibits discrimination on the basis of creed, sexual orientation, gender identity, religion, pregnancy or public accommodation. If you believe that you have been discriminated against in any program, activity or facility, as described above, or for more information, write Director, DNR, 502 E 9<sup>th</sup> St, Des Moines, IA 50319-0034 or the USDA, Director, Office of Civil Rights, 1400 Independence Ave SW, Washington, DC 20250-9410, or call (800) 795-3272 (voice) or (202) 720-6382 (TDD).

# Iowa's Forest Health Highlights

## Introduction

Each year the Iowa DNR Forestry cooperates with numerous agencies to protect Iowa's forests from insects, diseases, and other damaging agents. These programs involve ground and aerial surveys, setting up pheromone traps, following transects for sampling, collecting samples for laboratory analysis, and directing treatments for specific problems during the growing season. After each growing season, the Forestry Section issues a summary report regarding the health of Iowa's forests

This year's report begins with a brief summary of weather events, Iowa's land characteristics, and several survey summaries for insects, diseases, and invasive plants that have the potential to impact the health of Iowa's forests. The 2022 Forest Health Highlights will focus first on the Forest Service's Major Forest Pest List (Page 6) and then cover the additional damaging agents that DNR surveyed.

## Weather Review

This winter brought about average temperatures and on average levels of precipitation. There were several days in January that went above freezing, which caused many conifers to break winter dormancy. The repeated breaks in winter dormancy allowed for winter desiccation and eventual tree death in many conifer species throughout the state. Ongoing statewide severe drought conditions and extreme wind events in the west also contributed to conifer decline.


The entire state experienced cooler than average spring temperatures with most all of Iowa receiving fewer rainfall events. Anthracnose (a fungal leaf disease) on sycamore and many other benign fungal leaf diseases were common throughout the state despite the drier conditions. Tubakia and other leaf spot fungi were reported on oaks throughout the summer months and the number of reports increased slightly over last season.

Most of the state experienced normal summer temperatures statewide. However, drought conditions were present in most of the state with the driest condition occurring in central Iowa. Numerous reports of bur oak blight were made, in addition to frequent benign fungal leaf diseases that were being reported, in Eastern Iowa. DNR document an increase of 28% of oak leaf disease in 2022. Only bur oak blight was of concern.


DNR and partners have added 9 new counties of positive cases for the Emerald Ash Borer (EAB) in 2022. There are 93 of the 99 counties that have EAB confirmed. Several phone calls and emails were received about the increased level of ash mortality. Although there are many replanting efforts currently in place to offset this loss, DNR continues to provide resources to those communities who have been hit the hardest by this pest as funding allows.

Efforts have also been made to survey and identify over 7,800 acres of damage of white oak mortality (WOM). DNR has seen increased cases of WOM within the southeastern part of Iowa. This complex is believed to be linked to a number of factors including: Armillaria root disease, two-lined chestnut borer, drought, oak wilt, and other climate related issues. DNR continues to monitor the WOM locations. New survey methods are being implemented for the 2023 season to help pinpoint the cause and is expected to be reported in the 2023 Forest Health Highlights.


Average Temperature (°F): Departure from Mean  
December 1, 2021 to February 28, 2022




Average Temperature (°F): Departure from Mean  
March 1, 2022 to May 31, 2022




Average Temperature (°F): Departure from Mean  
June 1, 2022 to August 31, 2022




Accumulated Precipitation: Percent of Mean  
December 1, 2021 to February 28, 2022



Accumulated Precipitation: Percent of Mean  
March 1, 2022 to May 31, 2022




Accumulated Precipitation: Percent of Mean  
June 1, 2022 to August 31, 2022



[Images provided by Midwest Climate Watch](#)

**U.S. Drought Monitor**  
**Iowa**



**October 18, 2022**  
(Released Thursday, Oct. 20, 2022)  
Valid 8 a.m. EDT

Drought Conditions (Percent Area)

	None	D0-D4	D1-D4	D2-D4	D3-D4	D4
<b>Current</b>	0.00	100.00	84.15	35.05	7.11	0.02
<b>Last Week</b> 10-11-2022	0.00	100.00	57.46	26.99	7.07	0.02
<b>3 Months Ago</b> 07-19-2022	52.97	47.03	15.26	7.34	2.47	0.00
<b>Start of Calendar Year</b> 01-04-2022	50.98	49.02	11.87	0.00	0.00	0.00
<b>Start of Water Year</b> 09-27-2022	20.90	79.10	45.05	22.25	5.07	0.02
<b>One Year Ago</b> 10-19-2021	22.97	77.03	45.06	14.05	0.00	0.00

**Intensity:**

- None
- D0 Abnormally Dry
- D1 Moderate Drought
- D2 Severe Drought
- D3 Extreme Drought
- D4 Exceptional Drought

The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. For more information on the Drought Monitor, go to <https://droughtmonitor.unl.edu/About.aspx>



**Author:**

Adam Hartman  
NOAA/NWS/NCEP/CPC



[droughtmonitor.unl.edu](https://droughtmonitor.unl.edu)

**U.S. Drought Monitor**  
**U.S. States and Puerto Rico**



**October 18, 2022**  
(Released Thursday, Oct. 20, 2022)  
Valid 8 a.m. EDT

Drought Conditions (Percent Area)

	None	D0-D4	D1-D4	D2-D4	D3-D4	D4
<b>Current</b>	31.14	68.86	49.71	28.82	12.27	2.55
<b>Last Week</b> 10-11-2022	31.52	68.48	46.43	27.71	12.11	2.47
<b>3 Months Ago</b> 07-19-2022	36.68	63.32	44.57	30.11	15.08	3.77
<b>Start of Calendar Year</b> 01-04-2022	39.91	60.09	45.82	29.35	10.33	1.30
<b>Start of Water Year</b> 09-27-2022	36.92	63.08	42.65	25.36	10.45	2.14
<b>One Year Ago</b> 10-19-2021	48.34	51.66	38.88	28.14	17.36	5.58

**Intensity:**

- None
- D0 Abnormally Dry
- D1 Moderate Drought
- D2 Severe Drought
- D3 Extreme Drought
- D4 Exceptional Drought

The Drought Monitor focuses on broad-scale conditions. Local conditions may vary. For more information on the Drought Monitor, go to <https://droughtmonitor.unl.edu/About.aspx>

**Author:**

Adam Hartman  
NOAA/NWS/NCEP/CPC




[droughtmonitor.unl.edu](https://droughtmonitor.unl.edu)

## Land Characteristics

Iowa has approximately 2.85 million acres of forested land representing a decrease from 3.1 million acres in 2012. Most of Iowa's forests are native hardwood with oak, hickory, maple, basswood, walnut, ash, elm, cottonwood, and many other hardwood species. Less than 3% of Iowa's forests are conifer forests. There are currently 1.06 million acres of oak forest in Iowa.

Nearly 95% of the Forest Inventory Analysis (FIA) plots found one or more invasive plants competing with natives. The data also showed that over half of trees in Iowa are the preferred tree species by the nonnative pest spongy moth. In addition, the FIA report found that the average annual tree growth has declined while the average annual tree mortality has increased. Much of Iowa's small forests and trees that were along fencerows were cleared to allow for more profitable row cropping. The FIA data also indicated that succession to shade tolerant hardwoods (maples/ironwood) replacing shade intolerant hardwoods (oak/hickory) is continuing. These are alarming forest health trends. (Miles, P.D. Wed Mar 25 20:46:53 MDT 2016. Forest Inventory EVALIDator web-application version 1.6.0.01. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northern Research Station.)

Currently, there are 186 businesses in Iowa which utilize the wood grown in Iowa's forests. The forest products industry contributes over \$3.9 billion each year to Iowa's economy, including over 18,000 jobs for Iowans (Analysis by E.M. (Ted) Bilek, Economist, USDA Forest Service, Forest Products Laboratory, Madison, WI). Additional details can be found on page 192 of [Iowa's Forest's Today](#).



## United States Forest Service Major Pests List

This is a national list. Pests highlighted in red (\*) do not pertain to Northeastern Area and do not need to be reported. The items in blue have no known impact in Iowa at this time.

### Non-Native Pests

Asian Longhorned Beetle  
Balsam Woolly Adelgid  
Beech Bark Disease  
Browntail Moth  
Butternut Canker  
Dogwood Anthracnose  
Emerald Ash Borer  
Goldspotted Oak Borer\*  
Spongy Moth  
Hemlock Woolly Adelgid  
Laurel Wilt  
Oak Wilt  
Sirex Woodwasp  
Sudden Oak Death  
Thousand Cankers Disease  
White Pine Blister Rust  
Winter Moth

### Native Pests

Armillaria Root Disease  
Aspen Leafminer  
Bur Oak Blight  
Douglas-Fir Beetle\*  
Douglas-fir Black Stain Root Disease\*  
Fir Engraver\*  
Forest Tent Caterpillar  
Fusiform Rust  
Heterobasidion Root Disease  
Jack Pine Budworm  
Jeffrey Pine Beetle\*  
Large Aspen Tortrix  
Mountain Pine Beetle\*  
Northern Spruce Engraver  
Pine Black Stain Root Disease  
Polyphagous Shot Hole Borer\*  
Port-Orford-Cedar Root Disease\*  
Southern Pine Beetle  
Spruce Beetle  
Spruce Budworm  
Subalpine Fir Mortality\*  
Western Five-Needle Pine Mortality\*  
Western Pine Beetle\*  
Western Spruce Budworm\*  
Yellow-Cedar Decline\*

**United States Forest Service Major Pests List:  
Armillaria Root Disease**

Year: 2022

State: Iowa

Forest Pest

Common Name: Armillaria Root Disease

Scientific Name: *Armillaria spp.*

Hosts: Hardwoods and Conifers

Setting: N/A

Counties: N/A

Survey Methods: Ground

Acres Affected: N/A

Narrative: Armillaria root disease is fairly common in Iowa. The crown symptoms consist of branch dieback and crown thinning. The fungus produces a mycelial fan in recently killed trees just underneath the inner bark that often have a strong “mushroom” odor. The most common signs are the rhizomorphs that are produced just under the bark, and sometimes just on the bark surface. The rhizomorphs look like “shoestring”, which is why this fungus is sometimes called the shoestring fungi.

If a landowner needs assistance with armillaria root disease, please contact Tivon Feeley (DNR Forest Health Program Leader) at 515-275-8453 or the ISU Plant Diagnostic Clinic at 515-294-0581. More information can be found on the [USDA website](#).



Figure 1. Armillaria rhizomorphs under the bark. (Image: Robert L. Anderson, USDA Forest Service, Bugwood.org).



**United States Forest Service Major Pests List:  
Asian long-horned beetle**

Year: 2022

State: Iowa

Forest Pest

Common Name: Asian long-horned beetle

Scientific Name: *Anoplophora glabripennis*

Hosts: Maple, horsechestnut/buckeye, willow, elm, birch, and sycamore

Setting: Urban

Counties: Adair, Allamakee, Benton, Boone, Buchanan, Chickasaw, Clarke, Clayton, Fayette, Franklin, Howard, Ida, Iowa, Jackson, Johnson, Jones, Kossuth, Linn, Louisa, Lyon, Madison, Mills, Mitchell, Montgomery, O'Brian, Osceola, Pocahontas, Polk, Sac, Scott, Story, Warren, Webster, Winneshiek, Wright

Survey Methods: Ground

Acres Affected: N/A

Narrative: State legislative funds allowed DNR to conduct seven community inventories looking for invasive pests including Asian long horned beetle. Maples that had advanced dieback, dime-sized exit holes, or no obvious reason for the decline (e.g., girdling roots, construction damage, or planting depth) were destructively examined for Asian long horned beetle. All of the maples surveyed were healthy and did not have any evidence of Asian long horned beetle. A total of 12,897 maple were part of this survey effort.

Asian long-horned beetle has not been identified in Iowa. It is expected that survey work will resume in 2023. DNR asks all citizens to assist in the future monitoring efforts of this pest.

If beetles are found (Figure 1.) contact Rhonda Santos (USDA Public Information Officer) at 508-852-8044 and Robin Pruisner (State Entomologist) at 515-725-1465. [Asian long-horned beetle](#) information can be found on the USDA website.



**Figure 2. Adult Asian long-horned Beetle (Image: Dennis Haugen, USDA Forest Service, Bugwood.org).**

**United States Forest Service Major Pests List:  
Bur Oak Blight**

Year 2022

State: Iowa

Forest Pest

Common Name: Bur Oak Blight

Scientific Name: *Tubakia iowensis*

Hosts: Bur oak and Swamp White Oak

Setting: Rural Forests, Nursery, and Urban

Counties: Statewide

Survey Methods: Aerial, Ground, General Observation, and Culturing

Acres Affected: Approximately 1,050 acres

Narrative: Bur oak blight has been recognized in Iowa for only the last 15 years. However, it is suspected that the fungus that causes the disease has probably been here much longer. Theories on why bur oak blight has increased include: a shift in climate temperatures, more frequent rain events, older mature trees might be more susceptible, and that trees are more susceptible on sites that have a history of grazing or construction.

The disease can be found in most counties in Iowa, causing severe decline and mortality. Spring chemical injections, as needed, with propiconazole (Alamo) seem to control bur oak blight. However, some chemical burning (phytotoxic effects of the chemical) does occur. This control method works well in urban settings.

Currently, control measures have not been identified for woodland trees. Severely declining bur oaks have been harvested (salvaged) before they die. The estimated acres affected reflect the approximate acres of woodland salvage cuts. This does not reflect the urban damage, which cannot be quantified at this time.

There have been a few reports confirmed by the ISU NPDN Clinic of bur oak blight of swamp white oak. These appear to be true swamp white oak and not hybrids. The damage tends to be some venial necrosis and does not appear to be as severe on swamp white oak as it is on bur oak. These positive swamp white oaks will be followed to determine if the disease progresses over the years as it does on bur oaks.

All samples bur oak blight should be sent to the ISU Plant Diagnostic Clinic, who can be reached at 515-294-0581.




Figure 3. Current map of known locations of bur oak blight. (Image: Created by DNR based on locations provided by Dr. Harrington, ISU).

**United States Forest Service Major Pests List:  
Butternut Canker**

Year 2022

State: Iowa

Forest Pest

Common Name: Butternut Canker

Scientific Name: *Ophiognomonia clavigignenti-juglandacearum*

Hosts: Butternut

Setting: Rural Forest

Counties: Statewide

Survey Methods: General Observation

Acres Affected: Eastern half of Iowa (Scattered throughout roughly 2 million acres)

Narrative: Butternut canker is found throughout Iowa, but is largely concentrated in the Eastern half of Iowa where butternuts occur. The disease is fatal to native non hybrid butternuts.

No formal survey work was conducted on butternut canker in 2022. No suspect samples were submitted to DNR. No damage was reported in 2022.

If a landowner needs assistance with [butternut canker](#), please contact Tivon Feeley (DNR Forest Health Program Leader) at 515-275-8453 or the ISU Plant Diagnostic Clinic at 515-294-0581.



**Figure 4. Examples of canker found on butternut trees (Image: Minnesota Department of Natural Resources Archive, Minnesota Department of Natural Resources, Bugwood.org).**

**United States Forest Service Major Pests List:  
Emerald Ash Borer**

Year 2022

State: Iowa

Forest Pest

Common Name: Emerald Ash Borer

Scientific Name: *Agrilus planipennis*

Hosts: All Ash (*Fraxinus*) species

Setting: Rural Forest, Nursery, Urban

Counties: Adair, Adams, Allamakee, Appanoose, Audubon, Benton, Black Hawk, Boone, Bremer, Buchanan, Buena Vista, Butler, Calhoun, Carroll, Cass, Cedar, Cerro Gordo, Cherokee, Chickasaw, Clarke, Clay, Clayton, Clinton, Crawford, Dallas, Davis, Decatur, Delaware, Des Moines, Dickinson, Dubuque, Fayette, Floyd, Franklin, Fremont, Greene, Grundy, Hamilton, Hancock, Hardin, Harrison, Henry, Howard, Humboldt, Ida, Iowa, Jackson, Jasper, Jefferson, Johnson, Jones, Keokuk, Kossuth, Lee, Louisa, Lucas, Lyon, Madison, Mahaska, Marion, Marshall, Mills, Mitchell, Monroe, Montgomery, Muscatine, O'Brien, Page, Pocahontas, Polk, Pottawattamie, Ringgold, Sac, Scott, Shelby, Sioux, Story, Tama, Taylor, Union, Van Buren, Wapello, Warren, Washington, Wayne, Webster, Winnebago, Winneshiek, Worth and Wright


Survey Methods: Aerial, Ground, General Observation, and Trapping

Acres Affected: 1,422,180 aerial acres

Narrative: Emerald ash borer (EAB) was identified and confirmed in Iowa on May 14, 2010 on Henderson Island in Allamakee County. EAB has since been confirmed in 93 counties. Since the insect was already widespread, a statewide quarantine was issued February 4, 2014.

DNR visually inspected ash trees in 9 counties in 2022. The surveys found EAB in Clay, Dickinson, Hancock, Humboldt, Ida, Kossuth, Mitchell, O'Brien, Sioux counties. The other counties were confirmed IDALS through inspections, reports made by arborist, municipal contacts, local citizens and County Conservation Boards.

If a landowner has an ash tree that they believe has emerald ash borer please contact Tivon Feeley (DNR Forest Health Program Leader) at 515-725-8453 or Mike Kintner (IDALS EAB Coordinator) at 515-725-2877. Emerald ash bore information can be found at [Emerald Ash Borer Info](#) and the [Iowa DNR](#).



**Figure 5. Locations of the emerald ash borer infestations, as of August 2022. Please note that the entire State of Iowa is now quarantined for EAB. The brown circles around each infestation represent a 15 miles radius. The brown circles are done to assist landowners that are considering chemical treatments. The current recommendations from the Iowa EAB Team are not to chemically treat an ash tree until your property is within one of the brown circles. (Image: Tivon Feeley, DNR)**

**United States Forest Service Major Pests List:  
Forest Tent Caterpillar**

Year 2022

State: Iowa

Forest Pest

Common Name: Forest Tent Caterpillar

Scientific Name: *Malacosoma disstria*

Hosts: Many tree species

Setting: Rural Forests and Urban

Counties: Allamakee, Chickasaw, Clayton, Delaware, Fayette, Howard, and Winneshiek

Survey Methods: Ground and General Observation

Acres Affected: Approximately 306 acres

Narrative: Iowa DNR started receiving reports of forest tent caterpillars in Northeast Iowa in late May. Forest tent caterpillars are native and commonly found throughout the United States. The forest tent caterpillars have regional outbreaks every 6 to 16 years.

This is the first year the numbers have started to increase of [forest tent caterpillars](#).



**Figure 6. The picture above shows forest tent caterpillars on the main stem of a young tree. (Image: Robert Honeywell, DNR).**

## United States Forest Service Major Pests List:

### *Lymantria dispar*

Year 2022

State: Iowa

Forest Pest

Common Name: Spongy Moth (Formerly Known as Gypsy Moth)

Scientific Name: *Lymantria dispar*

Hosts: Oak, spruce, maples, elms, and many more

Setting: Rural Forests and Urban

Counties: Statewide

Survey Methods: Pheromone Delta Traps

Acres Affected: None

Narrative: Spongy moth has repeatedly been captured in Iowa, but the population level has effectively been controlled through trapping and mating disruption. Feeding damage has not occurred to Iowa's trees.

During the 2022 trapping season, 752 male moths were captured. 28 areas will be 'delimit' trapped in 2023 to determine whether Iowa has an isolated early infestation, which adds an additional 303 traps. There are no treatment blocks identified for 2023

No egg masses were found during the fall survey. All maps will be final after genetic testing on the captured male moths is completed and shows that they are all spongy moth.

More information on the spongy moth and the spray program can be found on the [Iowa Tree Pest](#) website.




Slow the Spread efforts are coordinated by a chartered, non-profit foundation that coordinated the operations of the program and facilitates the movement of funding between federal (USDA Forest Service and APHIS) and state agencies. Currently there are 11 States that participate in STS, and a Board of Directors leads the foundation. The STS Foundation Board of Directors is responsible for managing the STS project, which includes approving the annual plan of work, and allocating available resources. Two groups located at Virginia Polytechnic Institute and State University (VT) and Michigan State University (MSU) manage the centralized database for the program. Ongoing research from VT and other researchers are helping to improve management options for *Lymantria dispar* and improve the modeling of *Lymantria dispar* populations. Together they have achieved their goal of reducing spread of this destructive pest by more than 60%, which has prevented infestation of more than 140 million acres in 15 years.

**The Threat:** *Lymantria dispar* is a destructive, exotic forest pest that feeds on over 300 species of trees. It was accidentally introduced into the United States in 1869 and is currently established throughout the northeast and parts of the upper mid-west (gray shaded area on maps), where it has defoliated 80 million acres since 1970.

- It feeds on over 300 species of trees but oaks are most preferred.
- Defoliation causes extensive tree mortality, reduces property values, adversely affects commerce and causes allergic reactions in sensitive individuals that come in contact with the caterpillars.
- Most of the susceptible hardwood forests in the United States are not yet infested and are still at risk.

**The Benefits:**

- Prevents invasion of more than 300 million acres over the next 30 years
- Protects the extensive urban and wild land hardwood forests in the south and upper mid-west while also protecting the environment through use of spongy moth specific strategies.
- Yields a benefit to cost ratio of 3 to 1 by delaying the onset of impacts that occur as spongy moth invades new areas. The 20-year net present value after subtracting costs ranges from 184 to 348 million dollars.
- Delays impacts associated with spongy moth quarantines on intra- and inter-state commerce.
- Unifies the partners and promotes a coordinated, region-wide action based on biological need.






Figure 7. The map above details the locations of all the spongy moth traps and the number of moths captured in them during the 2022-trapping season. The total male moth capture was 738 male moths. (Image: Mark Hollister, PPQ).

## United States Forest Service Major Pests List: Heterobasidion Root Disease

Year 2022

State: Iowa

Forest Pest

Common Name: Heterobasidion root disease

Scientific Name: *Heterobasidion spp.*

Hosts: Conifers (All)

Setting: N/A

Counties: Lucas and Van Buren

Survey Methods: N/A

Acres Affected: N/A

Narrative: Heterobasidion root disease has been identified in Iowa and is a pest that can occur throughout Iowa on pines or red cedar. Historically it has been reported on jack pine in Stephens State Forest. Survey work was conducted at Yellow River State Forest for Heterobasidion root disease. Heterobasidion root disease has not yet been identified at Yellow River State Forest. If a landowner suspects Heterobasidion root disease, please contact the ISU Plant Diagnostic Clinic at 515-294-0581. See the [USDA website](#) for more information.



Figure 8. Example of heterobasidion root disease. (Image: William Jacobi, Colorado State University, Bugwood.org)

## United States Forest Service Major Pests List: Oak Wilt

Year 2022

State: Iowa

Forest Pest

Common Name: Oak Wilt

Scientific Name: *Ceratocystis fagacearum*

Hosts: All Oak Species

Setting: Woodlands and Urban

Counties: Statewide

Survey Methods: Aerial and Ground

Acres Affected: 351 acres

Narrative: DNR did not take any oak wilt samples this year. The DNR laboratory was set up only for insect identification for the 2022 season.

If a landowner feels that they have discovered [oak wilt](#), please contact the ISU Plant Diagnostic Clinic at 515-294-0581.



**Figure 9. The map above details the counties in Iowa with confirmed oak wilt. Oak wilt may occur in the non-shaded counties, but has not been confirmed by the ISU Diagnostic Clinic. (Image: Quinn Chavez, USFS).**

## United States Forest Service Major Pests List: Sudden Oak Death

Year: 2022

State: Iowa

Forest Pest

Common Name: Sudden Oak Death

Scientific Name: *Phytophthora ramorum*

Hosts: All Oaks

Setting: Rural Forests, Nursery, and Urban

Counties: Statewide

Survey Methods: N/A


Acres Affected: N/A

Narrative: Iowa did receive numerous notices of “trace forward” of suspected sudden oak death in 2022, meaning that potentially infected plant material had been shipped to Iowa. The notices were received too late in the season and the plants had been sold. Iowa’s weather conditions make it extremely unlikely for sudden oak death to survive. Stream baiting, to test for sudden oak death was not conducted in 2022 and is not planned for 2023 at this time.

If a landowner suspects that their plants have [sudden oak death](#), please contact Tivon Feeley (DNR Forest Health Program Leader) at 515-725-8453 or Robin Pruisner (State Entomologist) at 515-725-1465.



Ooze bleeds from a canker on an infected oak.



Black zone lines are found under diseased bark in oak.

**Figure 10. Two examples of the oozing canker found on an infected tree. The black lines under the bark are also symptomatic of sudden oak death. (Images: Joseph O’Brien, USDA Forest Service Pest Alert, and Bugwood.org)**

## United States Forest Service Major Pests List: Thousand Cankers Disease

Year 2022

State: Iowa

Forest Pest

Common Name: Thousand Cankers Disease

Scientific Name: *Pityophthorus juglandis* and *Geosmithia morbida*

Hosts: Walnut

Setting: Rural Forests, Nursery, and Urban

Counties: Statewide

Survey Methods: Ground, General Observation, and Culturing

Acres Affected: None

Narrative A total of 631 walnut trees were selected for the 2022 walnut twig beetle survey. A Lindgren four funnel dry trap with the walnut twig beetle pheromone developed by ISCAA Technologies was placed in a declining walnut tree for each location in the survey.

The traps were left on the trees for three weeks before being moved to another tree during the months of April, May, and the first week in June.

The following beetles were collected during the survey: *Xyleborus atratus*, *Ambrosiodmus tachygraphus*, *Hylocurus rudis*, *Xylosandrus germanus*, *Xyleborinus saxeseni*, *Xyloterinus politus*, *Xylosandrus crassiusculus*, *Pityophthorus lautus* (and subspecies), *Pityophthorus crinalis*, and *Pityophthorus consimilis*. There was a total of 10,478 ambrosia beetles, Pityophthorus beetles, and weevils that were collected. **No walnut twig beetles were found during the survey.** There were numerous other beetles, not of concern, collected (i.e. Japanese beetle, June bugs, etc.) but not counted as part of the survey.

The highest beetle captures occurred during the months of May and June. Historically, the captures decreased after those months. These trends have been consistent over the last several years, indicating that the best time to capture *Pityophthorus* species in Iowa is May and the first week in June.

If a landowner has walnut trees that they believe have [thousand cankers disease](#), please contact the ISU Plant Diagnostic Clinic at 515-294-0581.



**Figure 11. One of the Lindgren funnel traps that were used in conjunction with the walnut twig beetle pheromone. The traps were placed at sawmills, communities, and campgrounds. (Image: Shane Donegan, DNR)**



**Figure 12. A look inside the Lindgren Funnel trap capture chamber. The picture shows two pheromone pouches and a 3 inch long strip of dog collar that was used to kill the beetles that entered the capture chamber. (Image: Shane Donegan, DNR)**





Figure 13. Microscopes were used to help identify the beetle captured. The walnut twig beetle is about 1/4 of an inch long.  
(Image: Shane Donegan, DNR)



Figure 14. Pictured above is a *Pityophthorus* sp. (not *P. juglandis*) that was captured and sent in for identification. (Image: Shane Donegan, DNR)





**Figure 15. The locations of the 631 survey traps for walnut twig beetle throughout the state. (Image: Tivon Feeley, DNR)**

## United States Forest Service Major Pests List: Blister Rust

Year: 2022

State: Iowa

Forest Pest

Common Name: White Pine Blister Rust

Scientific Name: *Cronartium ribicola*

Hosts: White Pine

Setting: N/A

Counties: N/A

Survey Methods: N/A

Acres Affected: Unknown

Narrative: White pine blister rust has been identified in Iowa, and is a pest that can occur throughout the native white pine range in Iowa. No additional funds were available to conduct survey work. No suspect samples were submitted to DNR or the ISU Plant Diagnostic Clinic. No other survey work was conducted for white pine blister rust. If a landowner suspects [white pine blister rust](#) they should contact the ISU Plant Diagnostic Clinic at 515-294-0581.




Figure 16. The range map for known areas of white pine blister rust (Map: USFS)



Figure 17. Rust spores on an infected tree. (Image: Brian Geils, USDA Forest Service, Bugwood.org)

**Additional Pest Surveyed:  
Spotted Lanternfly**

Year 2022

State: Iowa

Forest Pest

Common Name: Spotted Lanternfly

Scientific Name: *Lycorma delicatula*

Hosts: Maple, Oak, Pine, Poplar, Sycamore, Walnut, Willow

Setting: Nursery

Counties: Polk

Survey Methods: Ground observation

Acres Affected: Areas unknown

Narrative: Spotted Lanternfly was identified in Dallas county during the 2022 season. Two instars were found by staff at a nursery and later confirmed by IDNR, IDALS, and APHIS without a breeding population. DNR will continue its cooperative effort to help survey for this pest on 2023.

For more information about the spotted lanternfly More information on the spotted lanternfly can be found on the [Iowa DNR's Forest Health](#) website.



Figure 18. Adult Spotted Lanternfly (Image: Pennsylvania Department of Agriculture, Bugwood.org)



Figure 19. Immature (Instar) Spotted Lanternfly (Image: Lawrence Barringer, Pennsylvania Department of Agriculture, Bugwood.org)

**Additional Pest Surveyed:  
Cankerworm**

Year 2022

State: Iowa

Forest Pest

Common Name: Cankerworm, Loopers, Spanworms

Scientific Name: *Paleacrita vernata*

Hosts: Apple, Ash, Beech, Elm, Hickory, Linden, Maples and Oaks

Setting: Rural Forests and Urban

Counties: Benton, Cedar, Linn

Survey Methods: Ground observation

Acres Affected: Areas unknown

Narrative: Iowa DNR started receiving reports of cankerworm in select counties in late May of 2022. Cankerworm caterpillars are native and commonly found throughout the United States. The cankerworm caterpillar rarely causes defoliation that were observed in Benton, Cedar, and Linn counties. Expected that this pest will not be a risk to forest health. No formal survey work is planned to take place in 2023.



Figure 20. The picture above shows a Cankerworm on a leaf of a young tree. (Image: James B. Hanson, USDA Forest Service, Bugwood.org).



Figure 21. The picture above shows a tree that has been damaged by Cankerworm (Image: Tivon Feeley, DNR)

**Additional Pest Surveyed:  
Pine Shoot Beetle**

Year: 2022  
State: Iowa  
Forest Pest

Common Name: Pine Shoot Beetle  
Scientific Name: *Tomicus piniperda*

Hosts: All Pines  
Setting: Rural Forests, Nursery, and Urban  
Counties: Statewide  
Survey Methods: N/A  
Acres Affected: Unknown  
Narrative:

Pine Shoot Beetle was identified September 18, 2006 and all counties in Iowa were quarantined for pine shoot beetle. Since the entire state is quarantined, no further monitoring has been needed. If a landowner needs assistance with management options for the [pine shoot beetle](#), please contact the ISU Plant Diagnostic Clinic at 515-294-0581.




Figure 21. The map above shows the quarantined areas for pine shoot beetle. (Image: by USDA-APHIS-PPQ)

## Pine Shoot Beetle Background

The pine shoot beetle (*Tomicus piniperda* L.) is an introduced pest that attacks pines. It was first discovered in the US at a Christmas tree farm near Cleveland, Ohio, in July 1992. A native of Europe, the beetle attacks new shoots of pine trees, stunting the growth of the trees. The pine shoot beetle may also attack stressed pine trees by breeding under the bark at the base of the trees. The beetles can cause severe decline in the health of the trees, and in some cases, kill the trees when high populations of the beetle exist.

In May, 2006, USDA-APHIS-PPQ confirmed the presence of pine shoot beetle (PSB) in Dubuque and Scott counties. A Federal Order was issued effective June 22, 2006 placing Dubuque and Scott counties under a Federal quarantine for interstate movement of PSB regulated articles. Iowa Department of Agriculture and Land Stewardship (IDALS) was provided a copy of the Federal Order as well as additional information concerning the pine shoot beetle, and was requested to consider placing a state PSB quarantine for intrastate movement of PSB regulated articles from Dubuque and Scott Counties. However, after considerable review, IDALS declined to implement an intra-state quarantine for PSB. Therefore, a Federal Order was issued effective September 18, 2006 for quarantine of the entire state of Iowa for PSB, *Tomicus piniperda*.

The quarantine affects the following pine products, called “regulated articles”:

- Pine nursery stock
- Pine Christmas trees
- Wreaths and garlands
- Pine logs/lumber (with bark attached)

All pine nursery stock shipped from Iowa to a non-regulated state must be inspected and certified free from PSB. This inspection and certification must occur just before shipping. Small pine seedlings (less than 36 inches tall, and 1 inch in diameter) and greenhouse grown pines require a general inspection of the whole shipment. All other (larger) pine nursery stock shipments must have 100% tip-by-tip inspection.



**Figure 22.** The picture above shows the pine shoot beetle and the damage it causes to branches.  
(Images: Steve Passoa, USDA APHIS PPQ, Bugwood.org)

**Additional Pest Surveyed:  
Dutch Elm Disease**

Year 2022

State: Iowa

Forest Pest

Common Name: Dutch Elm Disease

Scientific Name: *Ophiostoma ulmi* or *Ophiostoma novo-ulmi*

Hosts: Elm

Setting: Rural Forests and Urban

Counties: Statewide

Survey Methods: Ground, General Observation, and Culturing

Acres Affected: Approximately 12,613 acres

Narrative: Dutch elm disease was introduced to North America in the 1930's and began killing millions of native elm trees. Dutch elm disease has been identified in all of Iowa's counties, and it's estimated that just over 95 percent of the urban elm trees have succumbed to this disease.

The fungus is native to Asia and was introduced to Europe shortly after World War I. From Europe, it traveled to North America in the 1930's in crates made from infected elm logs. The disease quickly infected elms across the United States since our native elms did not have natural resistance to the introduced pathogen.

Dutch elm disease was reported statewide in 2022. The 2022 season appeared to have a high occurrence of [Dutch elm disease](#).



**Figure 23. Areas where Dutch elm disease is generally known to occur within the continental United States.  
(Image: Tivon Feeley, DNR)**

**Additional Pest Surveyed:  
Hickory Dieback**

Year 2022

State: Iowa

Forest Pest

Common Name: Hickory Dieback

Scientific Name: *Fusarium solani* and *Ceratocystis smalleyi*

Hosts: Bitternut Hickory and Occasionally Shagbark Hickory

Setting: Rural Forests and Urban

Counties: Statewide

Survey Methods: Ground

Acres Affected: Approximately 12,604 acres

Narrative: Hickories have continued to decline statewide. Mortality has become fairly common within the range of bitternut hickory making it difficult to track and estimate the acres impacted. If a landowner suspects hickory mortality, they should contact the ISU Plant Diagnostic Clinic at 515-294-0581.



Figure 24. Hickory bark beetle attack. (Image: Dr. Jennifer Juzwik, USFS)



Figure 25. Associated cankers. (Image: Dr. Jennifer Juzwik, USFS)



## Additional Pest Surveyed:

### Invasive Plants

Exotic invasive species are plants that are non-native to an ecosystem and cause or are likely to cause economic or environmental harm to humans, crops, livestock, or natural plant and animal communities. The most common non-native species found in the FIA report as problematic in Iowa forests are multiflora rose, reed canary grass, bush honeysuckle, garlic mustard, Japanese knotweed, autumn olive, common buckthorn, Japanese barberry, and oriental bittersweet (Miles, P.D. Wed Mar 25 20:46:53 MDT 2016. [Forest Inventory DataMart](#) web-application version 1.6.0.01. St. Paul, MN: U.S. Department of Agriculture, Forest Service, Northern Research Station.).

These invasive and exotic plants are out-competing native forest species, diminishing fisheries and wildlife habitat, reducing water quality, reducing economic returns from forest management and tourism, and threatening long term forest sustainability and biodiversity. In 2013 Oriental bittersweet, Japanese knotweed, garlic mustard, and Japanese hops were made illegal to distribute in the State of Iowa.

### Known Invasive Plants in Iowa 2022

Key: NP= Not Present- Not known to exist in Iowa

I= Isolated- the species is infrequent, not commonly seen

LA= Locally Abundant- the species is present but is not in the majority of the counties

W= Widespread- commonly seen in the majority of counties in large or small populations

Species	Common Name	Abundance
<i>Abutilon theophrasti</i>	Velvetleaf	W
<i>Ailanthus altissima</i>	tree-of-heaven	W
<i>Alliaria petiolate</i>	garlic mustard	W
<i>Berberis thunbergii</i>	Japanese barberry	W
<i>Bromus tectorum</i>	cheatgrass	W
<i>Butomus umbellatus</i>	flowering rush	I
<i>Carduus acanthoides</i>	plumeless thistle	I
<i>Carduus nutans</i>	Musk thistle	W
<i>Celastrus orbiculata</i>	Oriental bittersweet	LA
<i>Centaurea maculosa/beibersteinii</i>	spotted knapweed	LA
<i>Centaurea repens</i>	Russian knapweed	I
<i>Centaurea solstitialis</i>	yellow starthistle	I
<i>Cirsium arvense</i>	Canada thistle	W
<i>Cirsium</i> spp.	thistle	W
<i>Cirsium vulgare</i>	bull thistle	W
<i>Conium maculatum</i>	poison hemlock	I
<i>Coronilla varia</i>	crown vetch	W
<i>Daucus carota</i>	Queen Anne's lace	W
<i>Dipsacus fullonum/sylvestris</i>	common teasel	I
<i>Dipsacus laciniatus</i>	cutleaf teasel	I
<i>Dipsacus sativus</i>	Indian teasel	NP
<i>Elaeagnus angustifolia</i>	Russian olive	I
<i>Elaeagnus umbellate</i>	autumn olive	LA
<i>Euonymus alatus</i>	burning bush	LA
<i>Euphorbia esula</i>	leafy spurge	W
<i>Fallopia japonica</i>	Japanese knotweed	LA
<i>Frangula alnus/Rhamnus frangula</i>	glossy buckthorn	I
<i>Heracleum mantegazzianum</i>	giant hogweed	NP

<b>Species</b>	<b>Common Name</b>	<b>Abundance</b>
<i>Hesperis matronalis</i>	dame's rocket	W
<i>Humulus japonicus</i>	Japanese hop	LA
<i>Lespedeza cuneata</i>	Sericea lespedeza	I
<i>Ligustrum japonicum</i>	Japanese privet	NP
<i>Ligustrum obtusifolium</i>	blunt-leaved or border privet	I
<i>Ligustrum sinense</i>	Chinese privet	NP
<i>Ligustrum vulgare</i>	common or European privet	I
<i>Lonicera fragrantissima</i>	fragrant honeysuckle	NP
<i>Lonicera japonica</i>	Japanese honeysuckle	LA
<i>Lonicera maackii</i>	Amur honeysuckle	W
<i>Lonicera standishii</i>	Standish's honeysuckle	NP
<i>Lonicera tatarica</i>	Tatarian honeysuckle	W
<i>Lonicera x bella</i>	Bell's honeysuckle	I
<i>Lonicera xylosteum</i>	European fly honeysuckle	NP
<i>Lythrum salicaria</i>	purple loosestrife	W
<i>Morus alba</i>	white mulberry	W
<i>Pastinaca sativa</i>	wild parsnip	W
<i>Potamogeton crispus</i>	curlyleaf pondweed	I
<i>Pueraria montana</i>	kudzu	I
<i>Rhamnus cathartica</i>	common buckthorn	W
<i>Rosa multiflora</i>	multiflora rose	W
<i>Tamarix</i> spp.	salt cedar	I




Figure 26. The map above details the locations of invasive species as identified by DNR District Foresters and the Forest Health Program Leader in 2022. (Image: Tivon Feeley, DNR)

**Additional Pest Surveyed:  
White Oak Mortality**

Year 2022

State: Iowa

Forest Pest

Common Name: White Oak Mortality

Scientific Name: Unknown

Hosts: Quercus alba

Setting: Rural Forests and Urban

Counties: SE Iowa

Survey Methods: General Observation

Acres Affected: Approximately 7,892 acres

Narrative: There have been several counties in SE Iowa where DNR has visited woodlands that had severe white oak mortality from unknown causes. The white oaks start to decline in the lower slopes and the decline/mortality quickly moves upland. The leaves turn chlorotic and within a year, the tree is complete dead. The current management plan is to aggressively harvest affected trees.

This pattern of decline is similar to what Missouri has reported over the past several years. Samples collected in 2017 indicated activity of Armillaria root disease, two-lined chestnut borer, and a variety of decline-inciting disease agents, so it is unclear of the foundational cause of mortality. Nested PCR tests for oak wilt disease have all been negative. In 2022, mortality continued, but the causes continue under investigation.

**Additional Pest Surveyed:  
Oak Tatters**

Year: 2022

State: Iowa

Forest Pest

Common Name: Oak Tatters

Scientific Name: Unknown

Hosts: Oaks and Hackberry

Setting: Rural Forests and Urban

Counties: Pottawatomie, Mills, Adair, Dallas, Polk, Hardin, Marshall, Jasper, Washington, Johnson, Linn, and Muscatine

Survey Methods: General Observation

Acres Affected: Unknown

Narrative: DNR received several phone calls of tatters on oak and hackberry in 2022. The cause of oak tatters is not known. However, a study conducted by the University of Illinois suggested that Class 5 herbicides might be causing oak tatters. This has yet to be proven in the field.

Overall, the number of reports received during 2022 were lower than average past growing seasons. DNR will continue to work with its cooperators to determine the cause of oak tatters.




Figure 27. The map below details the locations that reported oak tatters in 2022. (Image: Tivon Feeley, DNR)

## Conclusion

Management plays an important role in creating a healthy Iowa forest. The best insurance a person can have when managing their woodlands is diversity of tree species with the appropriate number of trees per acre. These simple management strategies may help prevent excessive tree loss from a single pest and help maintain the trees' vigor, which may make them more resistant to potentially destructive insects and diseases. The best management plan for community forests is to not have more than 10% of any one species represented. Iowa forests play an important role by providing abundant forest products and amenities, including outdoor recreation opportunities, wildlife habitat, water quality, human health, and the economic benefits of a vast array of wood and wood fiber products.

Iowa's forests are facing an unprecedented level of invasive pests, chemical damage, wildlife pressure, and improper management. Emerald ash borer, Spongy moth, bur oak blight, and thousand cankers disease on walnut could have a 91.6-billion-dollar impact on Iowa's woodlands and community trees. No longer will passive management allow for woodlands to be "preserved" in the condition that they are in today. Learning about your woodlands and how each component affects another will make it easier for Iowa's woodlands to be managed for long term health. If you need technical assistance with your woodlands contact your [private lands forester](#) for assistance.

The Forestry Section, through cooperation with other agencies, has programs in place to monitor forest stressors which have potential to move into Iowa and damage our forests. Those programs operated vigorously during 2022, and plans are in place for a similar continued vigorous forest health program operation in 2022. Those programs existed in part from funding received by USFS grants and the State of Iowa Woodland Health Appropriation.

However, budget constraints limit the amount of work for important matters such as: white oak decline, aspen decline, additional oak wilt pockets, and bur oak blight. Additional funds are needed for these important forest health issues to be addressed in 2022.

DNR would like to thank its collaborators from USDA-Forest Service, USDA-APHIS-PPQ, Iowa State University Extension, Iowa Department of Agriculture and Land Stewardship, and Department of Natural Resources Foresters.

**"A nation that destroys its soils destroys itself. Forests are the lungs of our land, purifying the air and giving fresh strength to our people."**

**- Franklin D. Roosevelt**

## Useful Phone Numbers and Websites

DNR Forestry Section has an updated [forest health page](#).

DNR maintains an [emerald ash borer resource page](#).

Iowa Department of Agriculture and Land Stewardship [Tree Health Page](#).

[Iowa State University's Pest Management](#) and the Environment page host information on emerald ash borer, Spongy moth, and much more.

The Iowa State University Plant Disease Clinic has been assisting Iowa for nearly 50 years and is still available to answer plant disease questions. From flowers to trees they are ready to help. Contact them at 515-294-0581 or check them out on their [Plant Disease Clinic website](#).

For the creepy and crawling things on your plants, don't forget to contact [Iowa State University Extension Entomology](#). They can help you identify the insect and discover the best control measures. Contact them 515-294-1101.

Check out the [DNR landowner assistance](#) web page.

Be sure to look at the updated [Iowa DNR website](#).

### DNR Forest Health Program Leader Contact Information

Tivon Feeley

Iowa Department of Natural Resources Forestry Bureau

502 E 9<sup>th</sup> St

Des Moines IA 50319

Phone: 515-725-8453

Fax: 515-725-8201

E-Mail: [tivon.feeley@dnr.iowa.gov](mailto:tivon.feeley@dnr.iowa.gov)