Martelle, IA



2011 Management Plan Prepared by Steven Swinconos Bureau of Forestry, Iowa DNR



Table of Contents

| Executive Summary | 4 |
|--|----------|
| Overview | 4 |
| Inventory and Results | 4 |
| Recommendations | 4 |
| Introduction | - |
| introduction | 5 |
| Inventory | 5 |
| | |
| Inventory Results | 6 |
| Annual Benefits | <i>6</i> |
| Annual Energy Benefits | |
| Annual Stormwater Benefits | |
| Annual Air Quality Benefits | |
| Annual Carbon Benefits | |
| Annual Aesthetics Benefits | |
| Financial Summary of all Benefits | 6 |
| Forest Structure | 7 |
| Species Distribution | 7 |
| Age Class | |
| Condition: Wood and Foliage | |
| Management Needs | |
| Canopy Cover | |
| Land Use and Location | |
| Recommendations | 8 |
| Risk Management | ۶ |
| Pruning Cycle | |
| Planting | |
| Continual Monitoring | |
| Six Year Maintenance Plan with No Additional Funding | 11 |
| Emerald Ash Borer | 4.4 |
| | |
| Ash Tree Removal | |
| EAB Quarantines | |
| Wood Disposal | |
| Canopy Replacement | |
| Postponed Work | |
| Monitoring Private Ash Trees | |
| Filvate Asii Tiees | 12 |
| | |
| Movilso Citad | 40 |
| Works Cited | 13 |
| Annendix A: i-Tree Data | 14 |

Executive Summary

Overview

This plan was developed to assist the City of Martelle with managing its urban forest, including budgeting and future planning. Trees can provide a multitude of benefits to the community, and sound management allows communities to best take advantage of these benefits. Management is especially important considering the serious threats posed by forest pests such as the emerald ash borer (EAB). EAB is an invasive insect imported from Eastern Asia on wood shipping crates that kills all species of ash trees (this does not include mountain ash). There is a strong possibility that 22% of Martelle's city owned trees (ash) will die once EAB becomes established in the community. With proper planning and management, the costs of removing dead and dying trees can be extended over years, mitigating public safety issues.

Inventory and Results

In 2010, a tree inventory was conducted using Global Positioning System (GPS) data collectors. The inventory was a complete inventory of street and park trees. Below are some key findings of the 18 trees inventoried.

- Martelle's trees provide \$2,350 of benefits annually, an average of \$131 a tree
- There are over 10 species of trees
- The top three groups are: Ash 22%, Red Oak 17%, and River Birch 11%
- 17% of trees are in need of some type of management
- 1 tree is recommended for removal

Recommendations

The core recommendations are detailed in the Recommendations Section. The Emerald Ash Borer Plan includes management recommendations as well. Below are some key recommendations.

- Of the 1 tree needing removal, it is 16 inches in diameter. *City ownership of the trees recommended for removal should be verified prior to any removal*
- 1 of the 3 ash trees are in need of follow up because they are displaying canopy die back which is a sign and symptom associated with EAB. It is not uncommon for ash trees to have canopy die back and sprouts on the tree, but the EAB is attracted to trees with these problems. The ash trees were examined and no EAB exit holes were found. Woodpecker feeding is an excellent way to look for EAB. Woodpeckers feed on emerald ash borer larvae located under the bark. Feeding is typically evident higher in the tree where the emerald ash borer prefers to attack first. Large numbers of larvae under the bark can lead to woodpecker damage that looks like strips of bark have been pulled off of the tree. This is called "flecking"
- All trees should be pruned on a routine schedule- one third of the city every other year
- Plant a diverse mix of trees that do not include: ash, maple, cottonwood, poplar, box elder, Siberian elm, evergreen, willow or black walnut
- Check ash trees with a visual survey yearly

Introduction

This plan was developed to assist Martelle with the management, budgeting and future planning of their urban forest. Across the state, forestry budgets continue to decrease with more and more of that money spent on tree removal. With the anticipated arrival of Emerald Ash Borer (EAB), an invasive pest that kills native ash trees, it is time to prepare for the increased costs of tree removal and replacement planting. With proper planning and management of the current canopy in City of Martelle, these costs can be extended over years and public safety issues from dead and dying ash trees mitigated.

Trees are an important component of City of Martelle's infrastructure and one of the greatest assets to the community. The benefits of trees are immense. Trees provide the community with improved air quality, stormwater runoff interception, energy conservation, lower traffic speeds, increased property values, reduced crime, improved mental health and create a desirable place to live, to name just a few benefits. It is essential that these benefits be maintained for the people of City of Martelle and future generations through good urban forestry management.

Good urban forestry management involves setting goals and developing management strategies to achieve these goals. An essential part of developing management strategies is a comprehensive public tree inventory. The inventory supplies information that will be used for maintenance, removal schedules, tree planting and budgeting. Basing actions on this information will help meet Martelle's urban forestry goals.

<u>Inventory</u>

In 2010, a tree inventory was conducted that included 100% of the city owned trees on both streets and parks. The tree data was collected using a handheld Global Positioning System (GPS) receiver. The data collector gives Geographic Information Systems (GIS) coordinates with an accuracy of 3 meters, which can be used in Arc GIS as an active GIS data layer. Because the inventory is a digital document the data can be updated with new information and become a working document.

The programming used to collect tree information on the data collectors was written to be compatible with a state-of-the-art software suite called i-Tree. i-Tree was developed by the USDA Forest Service to quantify the structure of community trees and the environmental services that trees provide. The i-Tree suite is a public domain which can be accessed for free.

To quantify the urban forest structure and benefits, specific data is collected for each tree. This data includes: location, land use, species, diameter at 4.5 ft, recommended maintenance, priority of that maintenance, leaf health, and wood condition. Additionally, signs and symptoms of EAB were noted for all ash trees. The signs and symptoms noted were canopy dieback, epicormic shoots, bark splitting, D-shaped borer exit holes, and wood pecker damage.

Inventory Results

The data collected for the 18 city trees was entered into the USDA Forest service program Street Tree Resource Analysis Tool for Urban forestry Management (STRATUM), part of the i-Tree suite. The following are results from the i-Tree STRATUM analysis. Findings

Annual Benefits

Annual Energy Benefits

Trees conserve energy by shading buildings and blocking winds. Martelle's trees reduce energy related costs by approximately \$737 annually (Appendix A, Table 1). These savings are both in Electricity (26 MWh) and in Natural Gas (3,617 Therms).

Annual Stormwater Benefits

City of Martelle's trees intercept about 34,435 gallons of rainfall or snow melt a year (Appendix A, Table 2). This interception provides \$933 of benefits to the city.

Annual Air Quality Benefits

Air quality is a persistent public health issue in Iowa. The urban forest improves air quality by removing pollutants, lowering air temperature, and reducing energy consumption, which in turn reduces emissions from power plants, and emitting volatile organic matter (ozone). In Martelle, it is estimated that trees remove 335 lbs of air pollution (ozone (O₃), particulate matter less than 10 microns (PM10), carbon monoxide (CO), nitrogen dioxide (NO₂), and sulfur dioxide (SO₂)) per year with a net value of \$128 (Appendix A, Table 3).

Annual Carbon Benefits

Carbon sequestration and storage reduce the amount of carbon in the atmosphere, mitigating climate change. In Martelle, trees sequester about 121,298 lbs of carbon a year with an associated value of \$910 (Appendix A, Table 5). In addition, the trees store 5,046 lbs of carbon, with a yearly benefit of \$76 (Appendix A, Table 4).

Annual Aesthetics Benefits

Social benefits of trees are hard to capture. The analysis does have a calculation for this area that includes: aesthetic value, property values, lowered rates of mental illness and crime, city livability and much more. Martelle receives \$475 in annual social benefits from trees (Appendix A, Table 6).

Financial Summary of all Benefits

According to the USDA Forest Service i-Tree STRATUM analysis, Martelle's trees provide \$2,350 of benefits annually. Benefits of individual trees vary based on size, species, health and location, but on average each of the 18 trees in Martelle provide approximately \$130 annually (Appendix A, Table 7).

Forest Structure

Species Distribution

Martelle has over 14 different tree species along city streets and parks (Appendix A, Figure 1). The distribution of trees by group is as follows:

| Maple | 3 | 17% (sugar, red, Norway) |
|---------------------|---|--------------------------|
| Ash | 4 | 22% (green, white) |
| Northern Red Oak | 3 | 16% |
| River Birch | 2 | 11% |
| Plum | 2 | 11% |
| Sweetgum | 1 | 6% |
| Broadleaf Deciduous | 2 | 11% |
| Blue Spruce | 1 | 6% |

Size Class

Most of Martelle's trees (42%) are between 18 and 24 inches in diameter at 4.5 ft (Appendix A, Figure 2). For size, a Bell Curve is preferred and shows the highest amount of trees around 18 - 24 inches in diameter at 4.5 ft. Martelle's size curve is in the middle, indicating an average size urban forest.

Condition: Wood and Foliage

Both wood condition and leaf condition are good indicators of the overall health of the urban forest. The foliage condition results for Martelle indicate that 89% of the trees are in good health, with 5% foliage in poor health, dead or dying (Appendix A, Figure 3 & Appendix B, Figure 3). Similarly, 78% of Martelle's trees are in good health for wood condition (appendix A, Figure 4 & Appendix B, Figure 3). Wood condition that is in poor health, dead or dying is about 5% of the population. This 5% is estimates of trees that need management follow up.

Management Needs

The following outlines the specific management needs of the street and park trees by number of trees and percent of canopy (Appendix B, Figure 3).

| None needed | 15 | 86% |
|-----------------|----|-----|
| Crown Cleaning | 0 | 10% |
| Crown Raising | 2 | 3% |
| Tree Staking | 0 | 0% |
| Tree Removal | 1 | 1% |
| Crown Reduction | 0 | 0% |

Canopy Cover

There is no canopy cover in Martelle (Appendix A, Figure 4). There is about 1 acre of canopy cover in the city park.

Land Use and Location

The majority of Martelle's city and park trees are in the park (Appendix A, Figure 6 & Appendix A, Figure7). The following describes the land use and locations for the street and park trees.

Land Use

Single family residential 11% Parks 89%

Location

Planting strip 11% Parks 89%

Recommendations

Risk Management

Hazardous trees can be a significant threat to both people and property. Trees that are dead or dying, or that have large issues such as trunk cracks longer than 18 inches should be removed. Broken branches and branches that interfere with motorist's vision of pedestrians, vehicles, traffic signs and signals, etc should be removed.

Hazardous trees

Martelle has 1 critical concern tree that need immediate removal. This tree can be seen on the Location of Trees with Recommended Maintenance map (Appendix B, Figure 4). It is recommended to start with the large diameter critical concern trees first. The 1 tree is between 12 and 18 inches in diameter at 4.5 ft. Please refer to the six year maintenance plan at the end of this section. After the critical concern tree is addressed, there should be follow up on the trees marked as mature trees intermediate concern. There are no trees with these needs. These trees should be evaluated within one year and maintenance work done.

Pruning Cycle

Proper pruning can extend the life and good health of trees, as well as reduce public safety issues. In the Management Needs section of the Findings there are four main maintenance issues to be addressed: routine pruning, crown cleaning, crown raising, and crown reduction. Crown cleaning removes dead, diseased, and damaged limbs. Crown raising is the removal of lower branches that are 2 inches in diameter or larger in the case of providing clearance for pedestrians or vehicles. Crown reduction is removing individual limbs from structures or utility wires. It is recommended that all trees be pruned on a routine schedule every five to seven years. Please refer to the six year maintenance plan for further information.

Pruning Practices


Below are some trees that have been pruned in a way that will make it difficult for these trees to respond correctly to the wounding. The good news is that this pruning can be easily corrected to allow for proper tree response.



Consider the guidelines when pruning:

- 1. To avoid concerns related to the fungus that causes the disease oak wilt, all oak species should only be pruned between October 1 and February 28th.
- 2. All final cuts should be outside the branch collar.
- 3. Unless pruning broken oak branches between March 1 and September 30th pruning paints are not needed.





Proper Pruning Cut

Planting

Most of the planting over the next 5 years will replace the trees that are removed. It is recommended to plant 1 to 2 trees for every tree removed, since survival rates will not be 100%. Please refer to the six year maintenance plan at the end of this section. It is not essential that the new trees be planted in the same location of the trees being removed. However, maintaining the same number of trees helps ensure continuation of the benefits of the existing urban forest in Martelle.

It is important to plant a diverse mix of species in the urban forest to maintain canopy health, since most insects and diseases target a genus (ash) or species (green ash) of trees. Current diversity recommendations advise that a genus (i.e. maple, oak) not make up more than 20% of the urban forest and a single species (i.e. silver maple, sugar maple, white oak, bur oak) not make up more than 10% of the total urban forest. Presently, there is a good mix of tree species (Appendix A, Figure 1). Also, ash trees have not been recommended since 2002, due to the threat of EAB. Other species to avoid because they are public nuisances include: cottonwood, poplar, box elder, Chinese & Siberian elm, evergreen as street trees, willow or black walnut.

Continual Monitoring

Due to the threat of EAB, it is important to continuously check the health of ash trees. It is recommended that ash trees be checked with a visual survey every year for tree death and for the following signs and symptoms: canopy dieback, epicormic shoots, bark splitting, D-shaped borer exit holes, and wood pecker damage.

Six Year Maintenance Plan

Year 1

Removal: 1 critical concern tree

Planting and Replacement: 1 tree to be planted in open locations

Inspection and Maintenance Mature trees: 11 trees Visual Survey for signs and symptoms of EAB

Year 2

Visual Survey for signs and symptoms of EAB

Year 3

Visual Survey for signs and symptoms of EAB

Year 4

Removal: As needed

Planting and Replacement: As needed

Routine trimming: As needed

Visual Survey for signs and symptoms of EAB

Year 5

Inspection and Maintenance Mature trees: 11 trees Visual Survey for signs and symptoms of EAB

Year 6

Routine trimming: As needed

Visual Survey for signs and symptoms of EAB

Emerald Ash Borer Plan

Ash Tree Removal

I do not recommend removal of healthy ash trees until the EAB is present in your community. *City ownership of the tree recommended for removal should be verified prior to any removal*

EAB Quarantines

EAB is an extremely destructive plant pest and it is responsible for the death and decline of over 25 million ash trees. Ash in both forested and urban settings constitute a significant portion of the canopy cover in the United States. Current tools to detect, control, suppress and eradicate this pest are not as robust as the USDA would desire. In order to stay ahead of this hard to detect beetle, the USDA is attempting to contain the beetle before it spreads beyond its known positions by regulating articles.

A regulated article under the USDA's quarantine includes any of the following items:

- emerald ash borer
- firewood of all hardwood species (for example ash, oak, maple and hickory)
- nursery stock and green lumber of ash
- any other ash material, whether living, dead, cut or fallen, including logs, stumps, roots, branches, as well as composted and not composted chips of the genus ash (Mountain ash is not included)

In addition, any other article, product or means of conveyance not listed above may be designated as a regulated article if a USDA inspector determines that it presents a risk of spreading EAB once quarantine is in effect for your county.

Wood Disposal

A very important aspect of planning is determining how wood infested with EAB will be handled, keeping in mind that quarantines will restrict its movement. Consider who will cut and haul the dead and dying trees? Is there an accessible, secured site big enough to store and sort the hundreds of trees and the associated brush and chips? How will wood be disposed of or utilized? Do you have equipment capable of handling the amount and size of ash trees your tree inventory has identified? Once your county is under quarantine for EAB, contact USDA-APHIS-PPQ at 515-251-4083 or visit the website

http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/regulatory.shtm I. Wood waste can be disposed of as you normally would if your county is not part of quarantine.

Canopy Replacement

As budget permits, all removed trees should be replaced. The new plantings should be a diverse mix and should **not** include ash, maple, cottonwood, poplar, box elder, Chinese & Siberian elm, evergreen, willow or black walnut as street trees.

Postponed Work

While finances, staffing and equipment are focused on the management of ash, usual services may be delayed. Tree removal requests on genus other than ash will be prioritized by hazardous or emergency situations only.

Monitoring

It is recommended that ash trees be checked with a visual survey every year for tree death and for the following signs and symptoms: canopy dieback, epicormic shoots, bark splitting, D-shaped borer exit holes, and wood pecker damage.

Works Cited

Census Bureau. 2000. http://censtats.census.gov/data/IA/1601964290.pdf (April, 2010)

USDA Forest Service, et al. 2006. i-Tree Software Suite v1.0 User's Manual. Pp. 27-40.

McPherson EG, Simpson JR, Peper PJ, Gardner SL, Vargas KE, Ho J, Maco S, Xiao Q. 2005b.

City of Charleston, South Carolina, municipal forest resource analysis. Internal Tech Rep. Davis, CA: U.S. Department of Agriculture, Center for Urban Forest Research. p. 57

Nowak, D.J. and J.F. Dwyer. 2007. Understanding the benefits and costs of urban forest ecosystems. In: Kuser, J. (ed.) Urban and Community Forestry in the Northeast. New York:

Springer. Pp. 25-46.

Peper, Paula J.; McPherson, E. Gregory; Simpson, James R.; Vargas, Kelaine E.; Xiao, Qingfu 2009. Lower Midwest community tree guide: benefits, costs, and strategic planting. Gen. Tech. Rep. PSW-GTR-219. Albany, CA: U.S. Department of Agriculture, Forest Service, Pacific Southwest Research Station. p.115

Appendix A: i-Tree Data



Table 1: Annual Energy Benefits

Annual Energy Benefits of Public Trees by Species

3/11/2011

| | Total Electricity | Electricity | Total Natural | Natural | Total Standar | % of Total | % of | Avg. |
|---------------------|-------------------|-------------|---------------|----------|---------------|------------|----------|---------|
| Species | (MWh) | (\$) | Gas (Therms) | Gas (\$) | (\$) d Error | Trees | Total \$ | \$/tree |
| Ash | 1.2 | 93 | 181.8 | 178 | 271 (N/A) | 22.2 | 36.8 | 67.80 |
| Northern red oak | 0.6 | 46 | 77.2 | 76 | 122 (N/A) | 16.7 | 16.5 | 40.57 |
| Broadleaf Deciduous | s 0.6 | 42 | 76.9 | 75 | 118 (N/A) | 11.1 | 16.0 | 58.81 |
| River birch | 0.6 | 44 | 87.0 | 85 | 130 (N/A) | 11.1 | 17.6 | 64.76 |
| Plum | 0.0 | 2 | 4.4 | 4 | 6 (N/A) | 11.1 | 0.9 | 3.13 |
| Norway maple | 0.0 | 0 | 0.8 | 1 | 1 (N/A) | 5.6 | 0.2 | 1.10 |
| Red maple | 0.0 | 0 | 0.7 | 1 | 1 (N/A) | 5.6 | 0.1 | 1.03 |
| Sugar maple | 0.3 | 24 | 44.2 | 43 | 68 (N/A) | 5.6 | 9.2 | 67.52 |
| Sweetgum | 0.0 | 2 | 3.7 | 4 | 6 (N/A) | 5.6 | 0.8 | 5.82 |
| Blue spruce | 0.1 | 5 | 10.2 | 10 | 15 (N/A) | 5.6 | 2.0 | 14.80 |
| Other street trees | 0.0 | 0 | 0.0 | 0 | 0 (N/A) | 0.0 | 0.0 | 0.00 |
| Citywide total | 3.4 | 259 | 487.0 | 477 | 737 (N/A) | 100.0 | 100.0 | 40.92 |

Table 2: Annual Stormwater Benefits

Annual Stormwater Benefits of Public Trees by Species

| Species | Total rainfall interception (Gal) | Total (\$) | Standard Error | % of Total Trees | % of Total \$ | Avg. \$/tree |
|--------------------|-----------------------------------|---------------|-------------------|---------------------|------------------|-----------------|
| Ash | 13,771 | 373 | (N/A) | 22.2 | 40.0 | 93.31 |
| Northern red oak | 4,425 | 120 | (N/A) | 16.7 | 12.9 | 39.98 |
| roadleaf Deciduous | 5,173 | 140 | (N/A) | 11.1 | 15.0 | 70.10 |
| iver birch | 6,243 | 169 | (N/A) | 11.1 | 18.1 | 84.60 |
| lum | 76 | 2 | (N/A) | 11.1 | 0.2 | 1.03 |
| orway maple | 12 | 0 | (N/A) | 5.6 | 0.0 | 0.33 |
| d maple | 12 | 0 | (N/A) | 5.6 | 0.0 | 0.32 |
| gar maple | 3,795 | 103 | (N/A) | 5.6 | 11.0 | 102.87 |
| reetgum | 172 | 5 | (N/A) | 5.6 | 0.5 | 4.65 |
| ue spruce | 755 | 20 | (N/A) | 5.6 | 2.2 | 20.47 |
| her street trees | 0 | 0 | (N/A) | 0.0 | 0.0 | 0.00 |
| ywide total | 34,435 | 933 | (N/A) | 100.0 | 100.0 | 51.85 |

Table 3: Annual Air Quality Benefits

Annual Air Quality Benefits of Public Trees by Species

711/2011

| | | De | position | (lb) | Total Avoided (lb) | | | | Total BVOC BVOC | | | | Total Standard % of Total Avg. | | | |
|---------------------|-----|--------|---------------------|---------|--------------------|--------|-----------------|-----|--------------------|-------------------|---------------------|------------------|--------------------------------|------------|-------|-------|
| Species | 03 | NO_2 | $^{\text{PM}}_{10}$ | so $_2$ | Depos. (\$) | NO_2 | ${\rm PM}_{10}$ | VOC | so ₂ Av | voided Er (\$) | nissions En (lb) | nissions (\$) | Total (lb) | (\$) Error | Trees | |
| Ash | 3.1 | 0.5 | 1.5 | 0.1 | 17 | 6.0 | 0.9 | 0.8 | 5.6 | 37 | -0.7 | -3 | 17.8 | 51 (N/A) | 22.2 | 12.73 |
| Northern red oak | 0.8 | 0.1 | 0.4 | 0.0 | 5 | 2.8 | 0.4 | 0.4 | 2.7 | 18 | -1.2 | -4 | 6.7 | 18 (N/A) | 16.7 | 6.02 |
| Broadleaf Deciduous | 1.1 | 0.2 | 0.5 | 0.0 | 6 | 2.7 | 0.4 | 0.4 | 2.5 | 17 | -0.3 | -1 | 7.5 | 21 (N/A) | 11.1 | 10.75 |
| River birch | 1.4 | 0.2 | 0.7 | 0.1 | 7 | 2.9 | 0.4 | 0.4 | 2.6 | 18 | -0.3 | -1 | 8.3 | 24 (N/A) | 11.1 | 11.87 |
| Plum | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.1 | 0.0 | 0.0 | 0.1 | 1 | 0.0 | 0 | 0.3 | 1 (N/A) | 11.1 | 0.41 |
| Norway maple | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 (N/A) | 5.6 | 0.14 |
| Red maple | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 (N/A) | 5.6 | 0.13 |
| Sugar maple | 0.5 | 0.1 | 0.2 | 0.0 | 3 | 1.5 | 0.2 | 0.2 | 1.4 | 9 | -0.4 | -1 | 3.9 | 11 (N/A) | 5.6 | 10.75 |
| Sweetgum | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.1 | 0.0 | 0.0 | 0.1 | 1 | 0.0 | 0 | 0.3 | 1 (N/A) | 5.6 | 0.87 |
| Blue spruce | 0.1 | 0.0 | 0.1 | 0.0 | 0 | 0.3 | 0.0 | 0.0 | 0.3 | 2 | -0.2 | -1 | 0.6 | 2 (N/A) | 5.6 | 1.53 |
| Other street trees | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0.0 | 0.0 | 0.0 | 0 | 0.0 | 0 | 0.0 | 0 (N/A) | 0.0 | 0.00 |
| Citywide total | 6.9 | 1.2 | 3.4 | 0.3 | 38 | 16.5 | 2.4 | 2.3 | 15.5 | 102 | -3.0 | -11 | 45.5 | 128 (N/A) | 100.0 | 7.14 |

Table 4: Annual Carbon Stored

Stored CO2 Benefits of Public Trees by Species

| Species | Total Stored CO2 (lbs) | Total Standar (\$) d Error | % of Total Trees | % of Total \$ | Avg. \$/tree |
|--------------------|---------------------------|-------------------------------|---------------------|------------------|-----------------|
| Ash | 50,786 | 381 (N/A) | 22.2 | 41.9 | 95.22 |
| Northern red oak | 15,408 | 116 (N/A) | 16.7 | 12.7 | 38.52 |
| Broadleaf | 17,904 | 134 (N/A) | 11.1 | 14.8 | 67.14 |
| River birch | 22,225 | 167 (N/A) | 11.1 | 18.3 | 83.35 |
| Plum | 192 | 1 (N/A) | 11.1 | 0.2 | 0.72 |
| Norway maple | 17 | 0 (N/A) | 5.6 | 0.0 | 0.13 |
| Red maple | 17 | 0 (N/A) | 5.6 | 0.0 | 0.13 |
| Sugar maple | 14,280 | 107 (N/A) | 5.6 | 11.8 | 107.10 |
| Sweetgum | 185 | 1 (N/A) | 5.6 | 0.2 | 1.39 |
| Blue spruce | 284 | 2 (N/A) | 5.6 | 0.2 | 2.13 |
| Other street trees | 0 | 0 (N/A) | 0.0 | 0.0 | 0.00 |
| Citywide total | 121,298 | 910 (N/A) | 100.0 | 100.0 | 50.54 |

Table 5: Annual Carbon Sequestered

Annual CO₂ Benefits of Public Trees by Species

3/11/2011

| Species | Sequestered (lb) | Sequestered (\$) | Decomposition Release (lb) | | Total Released (\$) | Avoided (lb) | Avoided (\$) | Net Total (lb) | Total Standar (\$) d Error | % of Total Trees | % of Total \$ | Avg. \$/tree |
|---------------------|---------------------|---------------------|-------------------------------|----|------------------------|-----------------|-----------------|-------------------|-------------------------------|---------------------|------------------|-----------------|
| Ash | 1,580 | | \ / | -1 | -2 | 2.056 | 15 | 3.391 | 25 (N/A) | 22.2 | 33.3 | 6.36 |
| Northern red oak | 945 | | -74 | -1 | -1 | 1.017 | 8 | 1,887 | 14 (N/A) | 16.7 | 18.5 | 4.72 |
| Broadleaf Deciduous | 756 | 6 | -86 | 0 | -1 | 934 | 7 | 1,603 | 12 (N/A) | 11.1 | 15.7 | 6.01 |
| River birch | 840 | 6 | -107 | 0 | -1 | 979 | 7 | 1,711 | 13 (N/A) | 11.1 | 16.8 | 6.42 |
| Plum | 47 | 0 | -1 | 0 | 0 | 43 | 0 | 88 | 1 (N/A) | 11.1 | 0.9 | 0.33 |
| Norway maple | 5 | 0 | 0 | 0 | 0 | 7 | 0 | 12 | 0 (N/A) | 5.6 | 0.1 | 0.09 |
| Red maple | 3 | 0 | 0 | 0 | 0 | 7 | 0 | 9 | 0 (N/A) | 5.6 | 0.1 | 0.07 |
| Sugar maple | 758 | 6 | -69 | 0 | -1 | 535 | 4 | 1,224 | 9 (N/A) | 5.6 | 12.0 | 9.18 |
| Sweetgum | 74 | 1 | -1 | 0 | 0 | 49 | 0 | 122 | 1 (N/A) | 5.6 | 1.2 | 0.91 |
| Blue spruce | 39 | 0 | -1 | 0 | 0 | 106 | 1 | 143 | 1 (N/A) | 5.6 | 1.4 | 1.07 |
| Other street trees | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 (N/A) | 0.0 | 0.0 | 0.00 |
| Citywide total | 5,046 | 38 | -582 | -4 | -4 | 5,732 | 43 | 10,192 | 76 (N/A) | 100.0 | 100.0 | 4.25 |

Table 6: Annual Social and Aesthetic Benefits

Annual Aesthetic/Other Benefits of Public Trees by Species


| | | Standar | % of Total | % of Total | Avg. |
|---------------------|------------|---------|------------|------------|---------|
| Species | Total (\$) | | Trees | \$ | \$/tree |
| Ash | 137 | (N/A) | 22.2 | 28.9 | 34.36 |
| Northern red oak | 76 | (N/A) | 16.7 | 15.9 | 25.21 |
| Broadleaf Deciduous | 71 | (N/A) | 11.1 | 14.9 | 35.31 |
| River birch | 75 | (N/A) | 11.1 | 15.7 | 37.26 |
| Plum | 2 | (N/A) | 11.1 | 0.4 | 1.05 |
| Norway maple | 3 | (N/A) | 5.6 | 0.6 | 2.74 |
| Red maple | 0 | (N/A) | 5.6 | 0.0 | 0.04 |
| Sugar maple | 76 | (N/A) | 5.6 | 16.1 | 76.42 |
| Sweetgum | 15 | (N/A) | 5.6 | 3.1 | 14.73 |
| Blue spruce | 21 | (N/A) | 5.6 | 4.4 | 21.08 |
| Other street trees | 0 | (±NaN) | 0.0 | 0.0 | 0.00 |
| Citywide total | 475 | (N/A) | 100.0 | 100.0 | 26.41 |

Table 7: Summary of Benefits in Dollars

Total Annual Benefits of Public Trees by Species (\$)

| Species | Energy | co ₂ | Air Quality | Stormwater | Aesthetic/Other | Total Standard (\$) Error | % of Total \$ |
|---------------------|--------|-----------------|-------------|------------|-----------------|------------------------------|------------------|
| Ash | 271 | 25 | 51 | 373 | 137 | 858 (±0) | 36.5 |
| Northern red oak | 122 | 14 | 18 | 120 | 76 | 350 (±0) | 14.9 |
| Broadleaf Deciduous | 118 | 12 | 21 | 140 | 71 | 362 (±0) | 15.4 |
| River birch | 130 | 13 | 24 | 169 | 75 | 410 (±0) | 17.4 |
| Plum | 6 | 1 | 1 | 2 | 2 | 12 (±0) | 0.5 |
| Norway maple | 1 | 0 | 0 | 0 | 3 | 4 (±0) | 0.2 |
| Red maple | 1 | 0 | 0 | 0 | 0 | 2 (±0) | 0.1 |
| Sugar maple | 68 | 9 | 11 | 103 | 76 | 267 (±0) | 11.4 |
| Sweetgum | 6 | 1 | 1 | 5 | 15 | 27 (±0) | 1.1 |
| Blue spruce | 15 | 1 | 2 | 20 | 21 | 59 (±0) | 2.5 |
| Other street trees | 0 | 0 | 0 | 0 | 0 | 0 (±0) | 0.0 |
| Citywide Total | 737 | 76 | 128 | 933 | 475 | 2,350 (±0) | 100.0 |


Species Distribution of Public Trees (%)



| Species | Percent | |
|---------------------|---------|--|
| Ash | 22.2 | |
| Northern red oak | 16.7 | |
| Broadleaf Deciduous | 11.1 | |
| River birch | 11.1 | |
| Plum | 11.1 | |
| Norway maple | 5.6 | |
| Red maple | 5.6 | |
| Sugar maple | 5.6 | |
| Sweetgum | 5.6 | |
| Blue spruce | 5.6 | |
| Other species | 0.0 | |
| Total | 100.0 | |

Figure 1: Species Distribution

Relative Age Distribution of Top 10 Public Tree Species (%)



| | DBH class (in) | | | | | | | | |
|---------------------|----------------|-------|-------|-------|-------|-------|-------|-------|-----|
| Species | 0-3 | 3-6 | 6-12 | 12-18 | 18-24 | 24-30 | 30-36 | 36-42 | >42 |
| Ash | 0.0 | 0.0 | 0.0 | 0.0 | 25.0 | 75.0 | 0.0 | 0.0 | 0.0 |
| Northern red oak | 0.0 | 0.0 | 0.0 | 66.7 | 33.3 | 0.0 | 0.0 | 0.0 | 0.0 |
| Broadleaf Deciduous | 0.0 | 0.0 | 0.0 | 50.0 | 0.0 | 50.0 | 0.0 | 0.0 | 0.0 |
| River birch | 0.0 | 0.0 | 0.0 | 0.0 | 50.0 | 50.0 | 0.0 | 0.0 | 0.0 |
| Plum | 50.0 | 50.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Norway maple | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Red maple | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Sugar maple | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 |
| Sweetgum | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Blue spruce | 0.0 | 0.0 | 100.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |
| Citywide total | 16.7 | 11.1 | 5.6 | 16.7 | 16.7 | 33.3 | 0.0 | 0.0 | 0.0 |

Figure 2: Relative Size Class

Functional (Foliage) Condition of Public Trees by Species (%)

3/11/2011




Figure 3: Foliage Condition

Structural (Woody) Condition of Public Trees by Species (%)





Figure 4: Wood Condition

Canopy Cover of Public Trees (Acres)




| Zone | Acres | % of Total Canopy Cover |
|----------------|-------|-------------------------|
| 1 | 0 | 100.0 |
| Citywide total | 0 | 100.0 |

| | Total Land | Total Street and Sidewalk | | Canopy Cover as % of Total Land | Canopy Cover as % of Total Streets and | | |
|----------|------------|------------------------------|-------|------------------------------------|---|--|--|
| | Area | | Cover | | Sidewalks | | |
| Citywide | 0 | 0 | 0 | | | | |

Figure 5: Canopy Cover in Acres


Land Use of Public Trees by Zone (%)



| Zone | Single family residential | Multi- family residential | Industrial/ Large commercial | Park/vacant/ other | Small commercial |
|----------------|---------------------------------|---------------------------------|------------------------------------|-----------------------|---------------------|
| 1 | 11.1 | 0.0 | 0.0 | 88.9 | 0.0 |
| Citywide total | 11.1 | 0.0 | 0.0 | 88.9 | 0.0 |

Figure 6: Land Use of city/park trees

Location of Public Trees by Zone (%)



| Zone | Front yard | Planting strip | Cutout | Median | Other maintained locations | Other un- maintained locations | Backyard |
|----------------|------------|-------------------|--------|--------|----------------------------------|--------------------------------------|----------|
| 1 | 0.0 | 11.1 | 0.0 | 0.0 | 88.9 | 0.0 | 0.0 |
| Citywide total | 0.0 | 11.1 | 0.0 | 0.0 | 88.9 | 0.0 | 0.0 |

Zone

Citywide total

Figure 7: Location of city/park trees

The State of Iowa is an Equal Opportunity Employer and provider of ADA services.

Federal law prohibits employment discrimination on the basis of race, color, age, religion, national origin, sex or disability. State law prohibits employment discrimination on the basis of race, color, creed, age, sex, sexual orientation, gender identity, national origin, religion, pregnancy, or disability. State law also prohibits public accommodation (such as access to services or physical facilities) discrimination on the basis of race, color, creed, religion, sex, sexual orientation, gender identity, religion, national origin, or disability. If you believe you have been discriminated against in any program, activity or facility as described above, or if you desire further information, please contact the Iowa Civil Rights Commission, 1-800-457-4416, or write to the Iowa Department of Natural Resources, Wallace State Office Bldg., 502 E. 9th St., Des Moines, IA 50319.

If you need accommodations because of disability to access the services of this Agency, please contact the Director at 515-281-5918.